



Molecular Biology and Inhibitors of Hepatitis A Virus

Journal:	<i>Medicinal Research Reviews</i>
Manuscript ID:	MED-13-010.R1
Wiley - Manuscript type:	Review Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Debing, Yannick; Rega Institute - University of Leuven, Microbiology and Immunology Neyts, Johan; Rega Institute - University of Leuven, Microbiology and Immunology Thibaut, Hendrik; Rega Institute - University of Leuven, Microbiology and Immunology
Keywords:	hepatitis A virus, virus replication, immune evasion, atopy, antiviral drugs

SCHOLARONE™
Manuscripts

View

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

MOLECULAR BIOLOGY AND INHIBITORS OF HEPATITIS A VIRUS

Yannick Debing, Johan Neyts* and Hendrik Jan Thibaut

Rega Institute for Medical Research, University of Leuven, Leuven, Belgium

Key words: hepatitis A virus, virus replication, immune evasion, atopy, antiviral drugs

* Corresponding author:
Johan Neyts
Rega Institute for Medical Research
Minderbroedersstraat 10
B-3000 Leuven
Belgium
Email: johan.neyts@rega.kuleuven.be
Fax: +32.(0)16.33.73.40
Phone: +32.(0)16.33.73.53

ABSTRACT

Hepatitis A virus (HAV) is a faeco-orally transmitted picornavirus and is one of the main causes of acute hepatitis worldwide. An overview of the molecular biology of HAV is presented with an emphasis on recent findings. Immune evasion strategies and a possible correlation between HAV and atopy are discussed as well. Despite the availability of efficient vaccines, antiviral drugs targeting HAV are required to treat severe cases of fulminant hepatitis, contain outbreaks and halt the potential spread of vaccine-escape variants. Additionally, such drugs could be used to shorten the period of illness and decrease associated economical costs. Several known inhibitors of HAV with various mechanisms of action will be discussed. Since none of these molecules is readily useable in the clinic and since the availability of an anti-HAV drug would be of clinical importance, increased efforts should be targeted towards discovery and development of such antivirals.

1
2
3
4 30 **1. Introduction**
5

6 31 Hepatitis A virus (HAV) is a major cause of enterically-transmitted hepatitis worldwide, posing a global
7
8 32 burden estimated in 2005 at 119 million infections of which 31 million resulted in symptomatic illness
9
10 33 and 34 000 in death.¹ The virus is transmitted faeco-orally, mainly through close contact with infected
11
12 34 individuals or by consumption of contaminated food and drinking water. The extreme environmental
13
14 35 stability of the HAV particle contributes significantly to its transmission.^{2,3} HAV epidemiology correlates
15
16 36 with poor hygiene and living conditions. Consequently, virtually every adult in developing countries is
17
18 37 seropositive due to childhood infection. On the contrary, in regions with improved hygiene standards,
19
20 38 infection is often postponed to later age. In addition, disease severity is generally age-dependent:
21
22 39 infections are usually mild or asymptomatic in young children, whereas at older age, hepatitis A
23
24 40 frequently presents with classic symptoms of acute hepatitis (e.g. jaundice, fatigue, general malaise,...)
25
26 41 and a higher incidence of fulminant hepatitis, which may require liver transplantation.⁴ Fulminant
27
28 42 hepatitis occurs especially in those aged over 50 for which mortality rates up to 5.4% have been
29
30 43 reported.⁵ Additionally, HAV superinfections in chronic liver disease patients (e.g. hepatitis B or C) are
31
32 44 believed to increase morbidity and mortality,⁶⁻⁸ although these findings are still subject to debate.⁹ Since
33
34 45 highly efficient vaccines that provide long-lasting immunity have become available, HAV mortality and
35
36 46 morbidity has decreased dramatically.^{8,10} However, occasional outbreaks of hepatitis A, sometimes
37
38 47 resulting in fatal outcomes, still occur in industrialized countries. For instance, in the last decade,
39
40 48 outbreaks have been linked to the consumption of contaminated green onions, semi-dried tomatoes,
41
42 49 seafood, such as raw oysters and sushi, and other foodstuffs.¹¹⁻¹⁴
43
44

45
46 50 As posited earlier,¹⁵ research interest in the molecular biology and pathogenesis of HAV has decreased
47
48 51 substantially since the availability of a safe and efficient vaccine. Nonetheless, HAV remains an intriguing
49
50 52 and poorly understood virus and hepatitis A is still a public health problem in many countries. Here, we
51
52
53
54
55
56
57
58
59
60

will review key aspects of the biology of HAV with an emphasis on recent findings and the unique characteristics of the virus. The potential clinical use of antivirals against HAV will be discussed as well.

2. HAV genome organization and replication cycle

HAHV is a non-enveloped, single-stranded RNA virus with a positive-sense genome. Despite the fact that HAV is classified within the family of the *Picornaviridae*, it exhibits quite some differences compared to other members of this family and is consequently the sole member of the genus Hepatovirus.¹⁶ Based on phylogenetic analysis of full length VP1 sequences, the single HAV serotype is divided into 6 genotypes (I-VI).¹⁷ Genotypes I to III are human and can be subdivided into subgenotypes A and B (and potentially subgenotype IC¹⁸), whereas genotypes IV to VI are of simian origin.¹⁷ The HAV genome (as shown in figure 1) is approximately 7.5kb in length and consists of a 5' untranslated region (UTR), a single open reading frame (ORF) and a 3'-UTR with a polyadenosine-tract. Like other picornaviruses, the HAV genome lacks a cap structure; instead a VPg protein (3B) is attached to the 5' end. In the 5'-UTR, six secondary structure domains can be found: domain I contains a hairpin structure, while the second domain comprises two stem loop structures followed by a polypyrimidine tract (pY1).¹⁹ The remaining domains form the type III internal ribosome entry site (IRES) allowing cap-independent translation of the viral genome. The single ORF comprises the structural genes (VP1 to VP4 (=P1) and 2A) and the non-structural genes (2B-C and 3A-D). The short 3'-UTR contains two stem loops and/or a pseudoknot structure which are, together with domains I and II and the polypyrimidine tract of the 5'-UTR, crucial for viral RNA synthesis.²⁰

2.1 Receptor binding and cell entry

In 1996, the HAV cellular receptor (HAVcr-1) was identified on African green monkey kidney cells (AGMK) as an attachment and probably functional receptor.²¹ Consequently, the human homolog (HuHAVcr-1) was identified and characterized as a human HAV receptor.²² HAVcr-1 is also known to be a

1
2
3 76 marker for acute ischemic kidney injury (in this context referred to as kidney injury molecule 1 or KIM-
4
5 77 1)²³ and a regulator of T-cell based immunity (in this context referred to as T-cell immunoglobulin and
6
7
8 78 mucin domain-containing protein 1 or TIM-1).²⁴ Nevertheless, since HAVcr-1 is also expressed on other
9
10 79 organs,²² it is likely that additional receptors are required for HAV attachment and entry. For instance, it
11
12 80 has been suggested that TIM-3 promotes HAV entry without being a functional receptor.²⁵ In addition,
13
14 81 HAV-specific IgA are reported to mediate infection of hepatocytes via the asialoglycoprotein receptor
15
16 82 (ASGPR).²⁶ This and the fact that IgA was also identified as a natural ligand for HAVcr-1²⁷ may explain
17
18 83 why (and how) IgA-coated HAV can enter the hepatocytes through both HAVcr-1 and ASGPR, thereby
19
20 84 promoting enterohepatic circulation and continuous (endogenous) reinfections of the liver.²⁸ This
21
22 85 phenomenon is thought to play an essential role in prolonged and relapsing cases of hepatitis A. Only
23
24 86 the emergence of avid IgG antibodies can break the cycle and eventually clear the infection.²⁸
25
26
27
28

29
30 87 **2.2 Uptake and uncoating**

31
32 88 Following receptor-mediated binding to the cell surface, the HAV particle is internalized. This has been
33
34 89 suggested to occur through receptor-mediated endocytosis since HAV infection can be inhibited by
35
36 90 blockers of endosomal acidification such as monensin, ammonium chloride and chloroquine.²⁹⁻³²
37
38 91 However, the precise mechanism remains unclear to date. Upon binding to its receptor the particle is
39
40 92 destabilized, thus initiating the uncoating process and releasing the single-stranded RNA into the host
41
42 93 cell cytoplasm.³³ Uncoating has been reported to be a slow and asynchronous process with a reported
43
44 94 duration of 4 to 10 hours, in contrast to under an hour for poliovirus.³⁴ It has been proposed that this
45
46 95 asynchronicity may be due to the fact that the HAV inoculum contains a heterogeneous mixture of
47
48 96 mature virions and provirions. These provirions still require a maturation cleavage (of VP0 into VP2 and
49
50 97 VP4, as discussed below) following entry, prior to uncoating, and might therefore uncoat more slowly
51
52 98 than mature virions.^{34,35} During the uncoating process, dense, non-infectious HAV uncoating
53
54 99 intermediates are formed. Intriguingly, these particles do not appear to have an altered sedimentation
55
56
57
58
59
60

profile, thereby differing from the typical picornaviral A particle.³⁶ Finally, both low pH and Ca^{2+} ions are reported to play an important role in HAV receptor binding, uncoating and during the maturation cleavage of provirions following virus entry (as described above).^{31,32,35,37} Upon completion of uncoating, the single-stranded RNA is released into the host cell by an unknown mechanism.

2.3 Translation

The viral RNA, being released into the cytoplasm, is then translated into a single polyprotein. Similar to other picornaviruses, HAV employs an IRES, located in the 5' UTR, to direct cap-independent translation of the viral genome using the host ribosomal machinery. Most picornaviruses affect the host cell protein synthesis to favor a more efficient translation of the viral mRNA. This process is mediated by a proteolytic cleavage of eukaryotic initiation factor 4G (eIF4G), thereby inducing a complete shut-down of capped mRNA translation.³⁸ Intriguingly, unlike for other picornaviruses, it is thought that HAV IRES depends on eIF4G, as part of an intact eIF4F complex.³⁹ The fact that HAV has to compete with an intact host cell machinery may also explain its poor replication in cell culture (see section 5). However, a recent publication by Redondo *et al.* suggested that HAV IRES-driven translation can occur without intact eIF4G and that another factor may be crucial for this translation.³⁸

Next, the translated polyprotein is processed co- and posttranslationally in a series of proteolytic cleavages into several functional precursor and mature proteins. The primary posttranslational cleavage occurs at the junction between P1-2A and 2B (instead of the junction between P1 and 2A as for other picornaviruses) (figure 1).⁴⁰ All other processing steps (except the VP1-2A junction, as discussed below) are mediated by 3C^{pro} and its functional precursor 3ABC^{pro} .⁴¹ This is different from other picornaviruses which may use L^{pro} , 2A, 3C^{pro} and the 3CD^{pro} precursor.⁴² Intriguingly, HAV protein 2A, unlike for other picornaviruses, has no proteolytic activity nor does it contain a ribosome-skipping sequence,⁴² but seems to be implicated in morphogenesis (see section 2.6).

2.4 Regulating the balance between translation and replication

Since translation of viral proteins and RNA replication are competing processes, they must be balanced properly to allow efficient viral replication.⁴³ In order to tip the balance from translation to replication, the poly(A)-binding protein (PABP), as part of the eIF4F complex, is cleaved by HAV 3C^{pro}. The N-terminal cleavage product of PABP was shown to have an improved RNA-binding capacity compared to uncleaved PABP and may act as a dominant negative for IRES-mediated translation, thus favoring viral RNA synthesis.⁴⁴ Additionally, proteolytic cleavage of the poly(rC)-binding protein PCBP2, which interacts with the pyrimidine-rich tract pY1 in the HAV 5'-UTR, may also be implicated in regulating the balance between translation and viral RNA synthesis.¹⁹ Other enzymes that bind to the HAV IRES are polypyrimidine tract binding protein (PTB), glyceraldehyde 3-phosphate dehydrogenase and La autoantigen that respectively enhance, suppress and suppress translation.⁴⁵⁻⁴⁷

2.5 Replication

As for most RNA viruses, replication of the viral genome takes places in replication complexes that consist of rearranged cellular membranes containing both viral and host proteins. Studies revealed important roles for 2BC precursor and 2C proteins in the membrane rearrangements forming the replication complexes. These complexes are described as a tubular vesicular network and were thought to be of endoplasmatic reticulum (ER) origin.^{48,49} However, a recent publication described the mitochondrial localization of the 3ABC precursor protein suggesting that HAV replication complexes may be derived from the outer mitochondrial membrane.⁵⁰ In line with this hypothesis, exchanging the mitochondria-targeting 3A transmembrane domain for the ER-targeting poliovirus 3A transmembrane domain resulted in loss of replication competence.⁵¹

Within the replication complexes, the viral genome is transcribed into antisense RNA which subsequently serves as a template for the production of new viral genomes. During this process, the 3D

protein functions as an RNA-dependent RNA polymerase. The 3B (VPg) protein serves as a starting point for primer-independent transcription and is covalently linked to the 5' genome end. A conserved replication element near the 5' end of the 3D^{pol}-coding sequence likely directs uridylylation of VPg by 3D^{pol}, in such way priming VPg for initiation of RNA replication.⁵² During replication, the 3A/3B junction remains uncleaved and the transmembrane protein 3A serves as an anchor tethering the growing HAV RNA strand and associated proteins to the membranes of the replication complex,^{42,53} as is the case for other picornaviruses.⁵⁴ It has also been reported that the 2C protein of HAV binds to the 3' end of the antisense RNA. In this way protein 2C may be implicated in anchoring the negative sense RNA template to the membranes of the replication complex.⁵⁵

2.6 Capsid assembly

Despite the fact that the multi-tiered capsid assembly process is only poorly understood, HAV differs at various steps from other picornaviruses. Following the initial cleavage at the 2A/2B junction by 3C^{pro} (as discussed above), the N-terminal part of protein 2A coordinates proper folding of the P1-2A precursor protein. This precursor protein is then processed by 3C^{pro} and its stable precursor 3ABC^{pro} into VP0 (consisting of VP4 and VP2), pX precursor (consisting of VP1 and 2A) and mature VP3 protein.^{42,56} These building blocks assemble into pentamers and subsequently associate with viral RNA to form preprovirions. In a final processing step, VP0 will be cleaved into VP4 and VP2 and precursor pX into VP1 and 2A, yielding mature virus particles with only VP1, VP2 and VP3 proteins. The processing of pX is mediated by a yet unknown cellular protease.⁵⁷ Although the cleavage can be performed by extracellular enzymes like factor Xa and trypsin,^{58,59} maturation can also be executed by the lysosomal proteinase cathepsin L. This may indicate a role for lysosomal proteinases in maturation cleavage and potential targeting of HAV provirions to the early lysosomes for maturation cleavage.⁵⁹ On the other hand, VP0 processing is thought to be a self-catalytic process that is dependent on the presence of encapsidated RNA.⁶⁰ The resulting VP4 is rather small compared to other picornaviruses and lacks an N-terminal

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

myristoylation signal.⁶¹ Note that the mature VP4 protein has never been identified in purified virus stocks and its role in capsid formation remains unclear.^{15,60}

2.7 Release

Hepatitis A virus is an enterically transmitted virus that replicates mostly in hepatocytes before it is excreted via the bile into the faeces. Despite the fact that the exact details of HAV release remain elusive, the mechanism seems to differ depending on the cell type infected and is thought to involve either a vesicle-mediated cellular protein transport pathway or specialized hepatocellular transport proteins involved in bile secretion.⁶² Blank and colleagues demonstrated that following infection of polarized human intestinal epithelial Caco-2 cells, release of progeny virus was largely restricted to the apical membrane.⁶² In this way, virus is secreted mainly into the intestinal lumen resulting in an amplification of the HAV inoculum in the intestines and thus an increased viral shedding and spreading of the virus. However, it is still unclear by which mechanism HAV reaches the blood stream. A role for transcytosis by M cells present in Peyer’s patches in the ileum has been suggested, a mechanism that was found for poliovirus and reovirus as well.⁶² Another study described that infection of polarized human hepatocytes occurred most efficiently via the basolateral plasma membrane, after which more than 95% of progeny virus was exported through the basolateral membrane (into the bloodstream), rather than through the apical membrane (into the bile channels).⁶³ This contrasts with the *in vivo* observation that bile and faecal titers are considerably higher than serum titers.^{15,64} Reuptake and transcytosis of progeny virus by the hepatocytes into the bile channels is suggested as the mechanism for enteric secretion.

2.8 Particle structure

HAV is a nonenveloped icosahedral particle of approximately 27nm in diameter.⁶⁵ The mature capsid is composed of 12 pentamers each consisting of 5 copies of VP1, VP2 and VP3. Unlike for other

1
2
3 193 picornaviruses, VP4 appears not to be present in mature HAV particles (see section 2.6). Attempts to
4
5 194 produce high-resolution images of HAV particles remained unsuccessful so far. A 2006 review¹⁵ showed
6
7
8 195 a medium-resolution image obtained by cryo-electron microscopy suggesting the absence of the well-
9
10 196 defined canyon surrounding the 5-fold axis, in this way differing from other picornaviruses. However,
11
12 197 these results have not been confirmed or published separately to date. HAV has a limited number of
13
14 198 neutralization antigenic sites. The immunodominant antigenic site is composed of VP1 and VP3 residues
15
16 199 and is conformation-dependent.⁶⁶ A second antigenic site is the glycoprotein A binding site by which HAV
17
18 200 can bind to erythrocytes, causing hemagglutination. This process is optimal at acidic pH, but is impaired
19
20 201 at physiological pH. This suggests that HAV has evolved to escape erythrocyte binding and consequent
21
22 202 clearance.^{67,68} Indeed, a mutant of this binding site displayed increased clearance from the blood and
23
24 203 lower overall fitness, suggesting evolutionary constraints and explaining the low level of antigenic
25
26 204 variability of the glycoprotein A binding site.⁶⁸

205 3. Immune evasion mechanisms

206 Hepatitis A is clinically characterized by a prolonged asymptomatic phase before clinical illness becomes
207 apparent and a very limited type I interferon response (only in week 1-2 after HAV challenge).⁶⁹ HAV
208 replication remains largely undetected by the immune system for several weeks after infection and
209 intrahepatic RNA was shown to persist for over 48 weeks in experimentally infected chimpanzees.⁶⁹ To
210 this end, HAV employs several strategies to evade the host immune response which are rather different
211 from those used by other picornaviruses.

212 3.1 Targeting MAVS and TRIF

213 One of these evasion tactics is to ablate the innate immunity's alarming system that induces IFN
214 expression. HAV was shown to inhibit double stranded RNA-induced IFN β gene expression⁷⁰ by
215 interfering with the retinoic acid-inducible gene I (RIG-I) and melanoma differentiation-associated gene

1
2
3 216 5 (MDA-5) signaling pathways (figure 2).⁷¹ RIG-I recognizes single-stranded 5'-triphosphate RNA, while
4
5 217 MDA-5 detects single- and double-stranded picornaviral RNA covalently linked to VPg.⁷² Both RIG-I and
6
7
8 218 MDA-5 use an adaptor protein called mitochondrial antiviral signaling protein (MAVS, also known as IPS-
9
10 219 1, VISA or Cardif) that is localized on the outer membrane of the mitochondria.⁵⁰ Upon activation by RIG-
11
12 220 I or MDA-5, MAVS recruits and activates TANK-binding kinase 1 (TBK1) and inhibitor of NF- κ B kinase ϵ
13
14 221 (IKK ϵ). TBK1 and IKK ϵ are both responsible for the phosphorylation of IFN regulatory factor 3 (IRF-3),
15
16 222 eventually leading to IRF-3 dimerization, nuclear translocation and induction of IFN β transcription. HAV
17
18 223 proteins 3ABC^{pro}⁵⁰ and 2B⁷² have been described to interfere with MAVS, thereby disrupting the innate
19
20 224 cellular antiviral defense mechanism. Precursor protein 3ABC is a stable polyprotein processing
21
22 225 intermediate that requires both the 3A and the 3C^{pro} domain for the cleavage and inactivation of MAVS.
23
24 226 The transmembrane domain of protein 3A ascertains mitochondrial localization while the 3C^{pro} catalytic
25
26 227 site performs the actual proteolytic cleavage of MAVS. Mere 3C^{pro}, lacking the 3A domain, is incapable to
27
28 228 perform this proteolysis.⁵⁰ In addition, it has been demonstrated that protein 2B suppressed both MAVS
29
30 229 functioning and the kinase activities of TBK1 and IKK ϵ , thus synergistically suppressing RIG-I/MDA-5
31
32 230 signaling, although the exact mechanism still remains to be elucidated.⁷²
33
34
35
36
37
38 231 Toll-like receptor 3 (TLR3) provides an additional recognition mechanism for double-stranded RNA
39
40 232 (dsRNA). Activation also results in IRF-3 phosphorylation and induction of IFN β transcription, but unlike
41
42 233 RIG-I and MDA-5, the downstream action of TLR3 is mediated through TRIF (Toll/IL-1 receptor domain-
43
44 234 containing adaptor inducing IFN β) which activates the TBK1 and IKK ϵ kinases. A recent study reported
45
46 235 the proteolytic cleavage of TRIF by the 3CD precursor.⁷³ In analogy with the specificity of
47
48 236 3ABC^{pro}-induced cleavage of MAVS, 3C^{pro} without the 3D domain is not capable of performing this
49
50 237 cleavage. This suggests that the 3D^{pol} sequence is required to modify the substrate specificity without
51
52 238 requiring the catalytic polymerase activity. In addition, proteolytic cleavage of TRIF cannot be performed
53
54 239 by the 3ABC intermediate.⁷³
55
56
57
58
59
60

Cleavage of TRIF and MAVS has been reported for other picornaviruses as well, e.g. coxsackievirus B3 (MAVS and TRIF)⁷⁴ and enterovirus 71 (TRIF)⁷⁵. Unlike for HAV, 3C^{pro} of both viruses appeared to be sufficient for proteolysis. Interestingly, both MAVS and TRIF are also cleaved by the HCV NS3/4A protease. Despite the fact that the HAV and HCV proteases differ in cleavage specificity and are phylogenetically unrelated, they target the same pair of adaptor proteins, which may be considered a remarkable example of convergent evolution.⁷³

3.2 Coding biases

A second immune evasion strategy employed by HAV can be found in its codon bias or non-uniform usage of codons during translation.⁷⁶ Due to the lack of mechanisms to induce cellular shut-off, HAV has to compete with host mRNA for translation (as described above). Within this context, HAV has strategically adopted a codon bias towards rare codons, resulting in a highly deoptimized coding usage.^{77,78} This phenomenon contributes considerably to the low replication efficiency of HAV in several ways and seems to serve multiple purposes: first, the deoptimized codon use may reflect a strategy by which rare tRNAs are preferred over abundant tRNAs thereby avoiding competition with cellular tRNAs in the absence of an adequate host cell translation shut off.⁷⁷ Second, due to a decreased translation and replication rate, the virus is capable of keeping the cellular amount of dsRNA to a minimum, thus escaping the host cell antiviral response.⁷⁸ In addition, CpG-containing codons are particularly suppressed in the HAV coding sequence.^{79,80} This particular codon deoptimization is proposed to be associated with the recognition of unmethylated CpG by the innate immunity as a pathogen marker, contrary to the methylated CpG pairs of the host. Also in this way, HAV can evade the attention of the immune system. Third, the use of rare codons, usually grouped in clusters, is important for translation kinetics as such cluster may lead to a transient translation stop, a phenomenon known as ribosome stalling.⁸¹ It has been suggested, that this transient stop may be an important prerequisite for proper folding of HAV structural proteins thereby contributing to the low antigenic variability and extreme

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

environmental stability of HAV.^{77,78,82} Consequently, when HAV is adapted to replicate *in vitro* under circumstances of chemically-induced cellular shut-off, the resulting strains (with a ‘re-deoptimized’ coding sequence) display reduced environmental stability.⁸³ Given the preference of HAV for reduced RNA replication and translation levels, it is now comprehensible why HAV would employ such inefficient type III IRES.^{84,85} a very efficient IRES would not be compatible with escaping innate immunity and with the many ribosome stalls during translation.⁸³

In conclusion, the coding biases found in the HAV genome may result from evolutionary pressures towards (i) environmental stability (and thus controlled translation kinetics to obtain a resilient viral particle) and (ii) evading the host immune system.⁷⁹ Additional insights into the mechanisms at action would prove very valuable with regards to our understanding of virus evolution and virus-host interplay.

3.3 Modulation of regulatory T cell activity

In addition to manipulation of the innate immunity, HAV also modifies the adaptive immunity through its receptor binding. HAVcr-1 is a phosphatidylserine (PtdSer) receptor and functions as a potent T-cell co-stimulatory molecule that regulates activation of and tolerance induction (thus suppressing autoimmune responses) in T cells.²⁴ A recent study reported that HAVcr-1 is constitutively expressed on regulatory T (Treg) cells.⁸⁶ The function of these specialized T cells is to limit the magnitude of the immune response to diverse pathogens, in such way avoiding a hyperactive immune response and subsequent collateral (tissue) damage.⁸⁷ Several micro-organisms activate Treg in order to limit inflammation and tissue injury, as is the case for chronic HBV and HCV infections.⁸⁷ Conversely, binding of HAV to its receptor inhibits Treg cell functioning.⁸⁶ In this way, the host is overwhelmed by anti-self-responses since Treg cells normally suppress auto-immune reactions, allowing HAV to escape the attention of the immune system. In addition, HAV inhibits the production of transforming growth factor

286 β by Treg cells, thereby blocking T-effector-mediated anti-HAV responses. Complementary, HAV
287 stimulates IL-22 production which limits immune-mediated liver damage.⁸⁶

288 Taken together, by inhibiting Treg function, HAV creates an environment that favors viral replication and
289 establishment of infection. These findings may also provide an explanation why HAV may elicit auto-
290 immune hepatitis and extrahepatic manifestations including autoimmune hemolytic anemia, even
291 though these complications are rare.⁸⁸

292 **4. A link between hepatitis A and atopy**

293 The hygiene hypothesis states that the uprising of atopic disease, including asthma, allergic rhinitis and
294 atopic dermatitis in industrialized countries is due to the increased hygiene.⁸⁹ As HAV serostatus can be
295 regarded as a hygiene marker, its correlation with atopy was studied. It has been reported by several
296 groups that HAV infection is negatively correlated with hay fever, asthma and other atopic diseases.^{89,90}
297 However, other studies were unable to replicate such effect.^{91,92} In addition, McIntire *et al* described
298 that previous HAV infection may protect against atopy in individuals carrying a 6 amino acid-insertion
299 (designated 157insMTTTPV) in the gene encoding HAVcr-1 (which is present in 46-64% of the
300 population).⁹³ However, more recent studies failed to fully replicate these results.^{94,95} In addition,
301 157insMTTTPV appears to increase the susceptibility to severe HAV infection.⁹⁶ A double mechanistic
302 basis was suggested to (partially) explain this observation: (i) HAV binds more efficiently to the
303 157insMTTTPV-carrying HAVcr-1, resulting in a more effective receptor and (ii) natural killer T cells
304 expressing this long form of HAVcr-1 were more cytolytic against HAV-infected hepatocytes.⁹⁶ Thus, the
305 157insMTTTPV may protect from atopy, but predispose to severe HAV infections.

306 In conclusion, a possible relationship between a previous HAV infection, protection from atopic disease
307 and HAVcr-1 polymorphisms is still subject of debate, but represents an interesting area of research. If
308 HAV truly has a protecting effect, several questions should be addressed: does infection need to occur

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

during childhood? Has vaccination a similar effect? Can this relationship be used in finding a therapeutic target for atopy? Insights into this relationship may prove crucial in combating the rise of atopic disease in industrialized countries.

5. Cell culture

Unlike for most picornaviruses, culturing wild-type (wt) HAV in cell culture has proven to be quite a challenge as these strains replicate only very marginally *in vitro* (e.g. reference 97). HAV can be adapted to growth in cell culture through serial passaging; a process that introduces cell culture-adapting mutations stepwise into the viral genome. Such mutations have been found in the P2 region (especially in 2B and 2C)^{98,99} and in the 5'-UTR,^{100,101} but also in other parts of the genome.⁹⁸ Intriguingly, these cell culture-adapting mutations result in HAV strains that are highly attenuated *in vivo* (e.g. references 99,100). A major advance in culturing HAV was the selection of a Huh7 cell line (designated Huh7-A-I) which allows genetically stable growth of wt HAV without the accumulation of cell culture-adapting mutations.¹⁰⁴

Although wt HAV does not induce a cytopathic effect (CPE) in cell culture, cytopathic strains may arise during prolonged serial passaging.^{105,106} CPE appears to be cell type-specific and is mostly found in monkey kidney cell lines like FRhK-4 and BS-C-1. These selected virus strains induce apoptosis through ribosomal RNA degradation by RNase L^{107,108} and consequent caspase activation.¹⁰⁹ Interestingly, it seems possible to induce CPE with cell culture-adapted non-cytopathic strains under specific conditions of lower temperature (<34°C) and decreased cell density, both in FRhK-4 and human A549 cells.¹¹⁰ Similarly, we observed CPE for the PA21 strain (genotype IIIa) in FRhK-4 cells under these conditions (unpublished results).

6. The need for antivirals?

Despite the availability of an efficient vaccine, an antiviral drug against HAV would be of great use. First, since HAV growth *in vitro* is rather limited, the production of vaccines remains a painstaking process which in part explains the high cost.¹⁵ This cost is particularly a problem in relatively poor regions with improved hygiene where infections occur at later age and consequently are more severe.⁴ Second, antivirals could shorten the period of illness and decrease symptoms and associated economic costs in infected unvaccinated patients. Early treatment of these infected unvaccinated persons may also prevent severe cases of fulminant hepatitis. Third, an antiviral could be a useful tool in rapidly containing epidemics. Lastly, the potential emergence of vaccine-escape variants has been reported recently;¹⁸ an anti-HAV drug could therefore be instrumental in halting the spread of such virus strains. In conclusion, there is certainly a need for antiviral drugs for hepatitis A, given that they are safe, efficacious and preferably cheap.

Research into HAV antiviral drug discovery was mostly performed during the late eighties and early nineties. Afterwards, research efforts and funding waned due to the introduction of vaccines,¹⁵ although some interesting work has been done on amantadine and 3C^{pro} inhibitors in the last decade. An overview of the most relevant inhibitors is presented here (also see table 1).

6.1 Interferon

In 1984, IFN β was already demonstrated to efficiently inhibit the replication of HAV in human embryo fibroblasts.¹¹¹ HAV-infected cells were completely cleared following 5 passages with 1000 IU/ml of IFN β . Later, another study reported an effective concentration 50 (EC₅₀, minimal concentration required to reduce virus replication by 50%) of 90 IU/mL and a selectivity index (SI = 50% cytotoxic concentration (CC₅₀)/EC₅₀) of more than 100 for IFN α -2a.¹¹² In addition to the *in vitro* activity, a case study describes the *in vivo* efficacy of IFN β in three patients with fulminant hepatitis A and one patient with severe

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

hepatic failure due to HAV. Following a treatment with 3 million units per day, liver functioning ameliorated and all 4 patients survived.¹¹³ However, in addition to its high price and parenteral administration, the clinical use of IFN is associated with severe side effects hampering usage in less developed regions.

6.2 Amantadine

Amantadine was initially thought to inhibit HAV by increasing the intravesicular pH. However, its antiviral effect required an extended incubation when compared to NH₄Cl and other lipophilic amines (e.g. methylamine, dansylcadaverine).²⁹ In line with these findings, inhibition of HAV IRES-dependent translation was later reported as a mechanism of action (MOA).¹¹⁴ A follow-up study confirmed the moderate activity of amantadine in cell-based assays.¹¹⁵ In addition, an increased effect was observed when amantadine was combined with IFN α or IL-29 (IFN λ 1).^{115,116} However, in addition to a fairly limited effect *in vitro* (EC₅₀ of 58 μ M and a SI around 5¹¹⁷), pharmacokinetic studies demonstrated maximum plasma concentrations (C_{max}) following a multiple dosage scheme well below the reported EC₅₀,¹¹⁸ limiting its use in a clinical context.

6.3 The curious case of guanidine HCl

For guanidine HCl, a protein 2C-targeting anti-enterovirus inhibitor, mixed results were reported. The earliest publication detected no inhibition at 1 and 2mM,¹¹⁹ but other groups found moderate inhibition (around 50%) following treatment with 3mM¹²⁰ or 2.5mM.¹⁰⁷ Cho and colleagues reported reversible inhibition at 2mM but also reported enhancement of replication at 0.1-1mM.¹²¹ Conversely to these results, Yi *et al* reported strong inhibition of viral replication at concentrations as low as 0.1mM.⁹⁹ Several factors may explain the inconsistency of these data: (i) mutations in the different viral genomes studied, specifically in the 2C region; (ii) the cell type used in the antiviral assay; (iii) the multiplicity of

infection (MOI), which was found to be an important determinant for the antiviral activity,¹⁰⁷ and (iv) the detection method employed.

6.4 Miscellaneous small molecule inhibitors of the HAV replication cycle

Only limited information is available on other compounds with anti-HAV activity. Early steps of the replication cycle can be blocked by iota-carrageenan, which probably functions as an inhibitor of virus attachment and receptor binding.¹²² Atropine, at 1.7mM, proved to moderately reduce HAV replication by targeting the early steps of the HAV replication cycle as well (attachment and possibly uncoating).¹²³ Due to toxicity and limited antiviral activity, the clinical potential of both compounds is low. For glycyrrhizin, an aqueous extract of the licorice root, an EC₅₀ of 325µM and a SI of 15 were reported.^{117,124} Subsequent studies suggested inhibition of membrane penetration as the putative MOA.¹²⁴ In addition to its anti-HAV activity, glycyrrhizin proved also to be active against a relatively large panel of other viruses, including HBV, HCV, HSV and HIV, with several proposed mechanisms of action, as reviewed by Fiore *et al.*¹²⁵ However, glycyrrhizin may cause pseudohyperaldosteronism due to its aldosterone-like effects. In addition, the compound is readily metabolized following oral and intravenous administration and the attained plasma concentrations are considerably lower than the reported EC₅₀,¹²⁶ hampering clinical use for treatment of HAV-infected patients.

Probably the most potent small molecules reported thus far are 4',6-dichloroflavan and 4',6-dichloroisoflavan (EC₅₀ values of 6nM).¹²⁷ Inhibition of cell entry or viral uncoating were posited as possible mechanisms of action for the antiviral activity against HAV. These drugs were also found to be effective against rhinoviruses and poliovirus *in vitro*.¹²⁷ However, 4',6-dichloroflavan failed to protect volunteers from experimental rhinovirus infection when administered both orally or intranasally,^{128,129} raising doubts concerning its *in vivo* efficacy against HAV as well.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Other early stage inhibitors include chlorpromazine and chloroquine.³¹ The former is thought to block endocytosis of non-clathrin-coated vesicles, while the latter inhibits endosome acidification.³¹ Monensin, another endosome acidification blocker, yielded minor to extensive inhibition of HAV replication in different studies.^{30,31,62}

In addition to early stage inhibitors, several other HAV inhibitors were identified to target different steps of the intracellular replication process including RNA synthesis, translation and capsid assembly. However, the information on their precise MOA is very limited. For instance, ribavirin was demonstrated to have a moderate effect at 100µM,¹³⁰ showing an EC₅₀ of 94µM and a selectivity index of around 4.¹¹⁷ Improved anti-HAV activity was reported for pyrazofurin, an orotidine 5'-monophosphate decarboxylase inhibitor, but further development was halted due to toxicity (interference with nucleotide metabolism).¹¹⁷ Protamine reduced HAV replication with 97% at 100µg/mL and was suggested to function as a RNA polymerase inhibitor.¹²³ In addition, inhibitors of oxidative phosphorylation and 2-deoxy-D-glucose, a glycolysis inhibitor, were identified as inhibitors of HAV replication by unknown mechanisms.³⁰ Other inhibitors include brefeldin A^{62,107} and cycloheximide,¹⁰⁷ inhibitors of ER-to-golgi-transport and protein synthesis respectively. A moderate anti-HAV activity was also reported for amphotericin B¹³¹ and methisoprinol,¹³² drugs used against fungal infections and as an immunomodulator in the treatment of subacute sclerosing panencephalitis respectively. However, no data are available on the MOA.

In recent years, several new anti-HAV compounds were synthesized, but the antiviral activity remained rather limited.^{133–135} Other papers reported highly potent inhibitors of HAV replication, but did not report toxicity data, making it impossible to judge whether or not these are selective inhibitors.^{136–138}

6.5 Rational design of 3C^{pro} inhibitors

420 Considerable efforts have been undertaken to rationally design and develop inhibitors for HAV and
421 human enterovirus 3C^{pro} and the related coronavirus 3C-like protease (3CL^{pro}).¹³⁹ These enzymes have a
422 topology comparable to that of the chymotrypsin-like serine proteases, but are in fact cysteine
423 proteases.¹³⁹ Consequently, the general research strategy is to introduce thiol-reactive groups into the
424 catalytic site. Several of such protease inhibitors have been developed over the years, both modified
425 peptides and non-peptidic analogs. Despite the fact that the modified peptides are easier to design (by
426 merely mimicking the natural protease substrate), non-peptidic and peptidomimetic inactivators are
427 preferred, since they can be modified in order to increase stability and cellular and gastro-intestinal
428 uptake, properties that are difficult to attain for peptides. A similar approach has already been applied
429 to rhino- and enterovirus 3C^{pro} and resulted in the molecule rupintrivir which was halted after
430 unsuccessful phase II clinical trials.¹⁴⁰

431 The first HAV 3C^{pro} inhibitors developed were modified peptides based on the acetyl-Leu-Ala-Ala
432 sequence, serving as competitors for the natural peptide substrate Leu-Arg-Thr.¹⁴¹ Based on these
433 findings, peptidyl aldehyde and monofluoromethylketone (FMK) inhibitors were developed.^{142,143}
434 Resolving the crystal structure of HAV 3C^{pro} in complex with a peptidyl FMK inhibitor indicated that the
435 inhibitor covalently binds the catalytic cysteine.¹⁴⁴ One of these peptidyl FMK inhibitors was also tested
436 *ex vivo* and was shown to reduce HAV replication by 96%.¹⁴³ However, cellular uptake appeared to be a
437 major problem. In addition, several azaglutamine derivatives were found to be irreversible peptidic
438 inhibitors.¹⁴⁵

439 On the other hand, several non-peptidic antagonists were reported as well. Asymmetric
440 azodicarboxamides were found to be irreversible inhibitors of HAV and HRV 3C^{pro}.¹⁴⁶ Serine and
441 threonine β -lactones also inhibited HAV 3C^{pro}.^{147,148} Other inhibitors include monophenyl
442 pseudoxazolones of glycine (IC₅₀ of 4-6 μ M)¹⁴⁹ and cathepsin K inhibitor-based keto-glutamines.¹⁵⁰ Lastly,

halopyridinyl esters were reported as the most potent non-peptide HAV 3C^{pro} inhibitors thus far with IC₅₀'s as low as 53nM.¹⁵¹

Although recent publications in this field seem to focus increasingly on HRV 3C^{pro} and coronavirus 3CL^{pro},¹⁵² it would be wise not to neglect possible anti-HAV activity for potential future (off-label) use of a marketed HRV or coronavirus protease inhibitor for severe HAV infections.

6.6 RNA interference

RNA interference (RNAi) has been suggested as a future treatment of severe HAV infections.¹⁵³ Small interfering RNA (siRNA) targeting non-structural genes can effectively block replication.^{153,154} Inhibition was also reported when targeting domains IIIC and V of the HAV IRES.¹⁵⁵ Three consecutive applications of siRNA over 14 days resulted in a 3 log₁₀ infectivity titer reduction in persistently infected cells. However, resistance emerged fairly quickly through mutation of the target site, necessitating combination of different siRNA's.¹⁵⁴ Once clinical RNAi therapy would be an established treatment for other (infectious) diseases, it would be interesting to further explore this for hepatitis A. However, such clinical applications of RNAi are currently still at the experimental stage.

6.7 Novel antiviral assays

We recently reported the development of three assays to screen for and identify small molecule inhibitors of HAV replication.¹⁵⁶ A CPE reduction assay based on the cell culture-adapted HM175/18f strain (genotype IB) is amendable to high-throughput screening. RT-qPCR-based virus yield assays for HM175/18f and genotype IIIA strain PA21 were developed as well, allowing for confirmation and further characterization of the antiviral activity of selected molecules. The known inhibitors IFN α and amantadine were used to validate these assays. Using these assays, three enterovirus inhibitors with different targets were evaluated for anti-HAV activity. Pleconaril, a known capsid binder,¹⁴⁰ yielded (not unexpectedly) a very limited activity; inhibition by rupintrivir, a 3C^{pro} inhibitor, proved to be strain-

dependent¹⁵⁶ and enviroxime, a direct inhibitor of phosphatidylinositol-4-kinase III β (PI4KIII β) that induces resistance mutations in enterovirus protein 3A,¹⁵⁷ was inactive in all 3 systems. Recently, PI4KIII β has been identified as an essential host factor in the replication strategy of enteroviruses. Following recruitment by enterovirus protein 3A to ER-derived organelle membranes, a phosphatidylinositol-4-phosphate (PI4P)-rich environment is created, promoting viral RNA replication. The observation that HAV protein 3A is targeted towards the outer mitochondrial membrane instead of the ER,⁵⁰ together with the observed lack of activity for enviroxime in 2 different cell lines,¹⁵⁶ suggests that PI4KIII β kinase activity does not play a role in HAV replication.

6.8 Small animal models

A small animal model would be an additional prerequisite for the development and validation of anti-HAV compounds. The host range of HAV is restricted to humans and several non-human primates, e.g. tamarins and chimpanzees.¹⁵⁸ Some efforts have been undertaken towards the development of small animal models, e.g. virus adaptation to mouse cell lines or infection of guinea pigs.^{159,160} However, so far no robust model for clinical hepatitis A is available. Very recently, it has been found that the chimeric SCID/Alb-uPA mouse model, which supports the engraftment and proliferation of transplanted human hepatocytes, is susceptible to HAV infection (Pang D, personal communication). Despite the fact that these results still need to be published, they could represent a major leap forward for anti-HAV research.

7. Conclusion

Despite the fact that hepatitis A is a vaccine-preventable disease and infections are usually mild, continued attention for this pathogen is warranted. Outbreaks and occasional cases of fatal fulminant hepatitis A still occur and as the age of infection shifts upwards, the severity of the infection increases as well. Morbidity and mortality may also increase in patients with chronic liver disease, e.g. due to HBV or

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

HCV. Additionally, a recent report warned for the potential emergence of vaccine-escape variants.¹⁸

Taken together, these reasons warrant continuation and expansion of the current vaccination programs, but also the development of antiviral drugs against HAV, even though this may not be particularly interesting from a commercial point of view. So far no potent or selective inhibitors have been reported.

We therefore propose to focus antiviral efforts on marketed drugs and antivirals that are currently in clinical development. Off-label use of such drug would be a much appreciated therapy. In fact, we are currently employing our CPE reduction assay for screening of a library of FDA approved/marketed drugs and with promising results so far. Also on the fundamental biology level, HAV remains a largely understudied virus, despite many interesting features. Further investigation could provide unique insights into pathogenesis, tissue tropism determinants and replication strategies for HAV and picornaviruses in general.

8. Acknowledgement

Yannick Debing would like to thank the Research Foundation – Flanders (FWO) of which he is a fellow.

This work is supported by the European Union Seventh Framework Program (FP7/2007-2013) under SILVER grant agreement n° 260644 and by KU Leuven, geconcerteerde onderzoeksactie (GOA/10/014).

9. References

1. Rein D. Modeling the global burden of hepatitis A virus infections in 1990 and 2005. Paper presented at: The Liver Meeting 2011. The 62nd Annual Meeting of the American Association for the Study of Liver Diseases; 2011 Nov 3-8; San Francisco, CA.
2. Siegl G, Weitz M, Kronauer G. Stability of hepatitis A virus. *Intervirology* 1984;22:218–226.
3. Abad FX, Pintó RM, Bosch A. Survival of enteric viruses on environmental fomites. *Appl Environ Microbiol* 1994;60:3704–3710.
4. FitzSimons D, Hendrickx G, Vorsters A, Van Damme P. Hepatitis A and E: update on prevention and epidemiology. *Vaccine* 2010;28:583–588.
5. WHO. The immunological basis for immunization series – module 18: hepatitis A. Geneva, Switzerland: WHO press; 2011. 39 p.
6. Keffe E. Hepatitis A in patients with chronic liver disease - severity of illness and prevention with vaccination. *J Viral Hepat* 2000;7 Suppl 1:15–17.
7. Vento S, Garofano T, Renzini C, Cainelli F, Casali F, Ghironzi G, e.a. Fulminant hepatitis associated with hepatitis A virus superinfection in patients with chronic hepatitis C. *N Engl J Med* 1998;338:286–290.
8. Vogt TM, Wise ME, Bell BP, Finelli L. Declining hepatitis A mortality in the United States during the era of hepatitis A vaccination. *J Infect Dis* 2008;197:1282–1288.
9. Shouval D. Hepatitis: New doubts about preventing HAV superinfection in chronic HCV. *Nat Rev Gastroenterol Hepatol* 2012;9:367–368.
10. Ott JJ, Irving G, Wiersma ST. Long-term protective effects of hepatitis A vaccines. A systematic review. *Vaccine* 2012;31:3–11.
11. Wheeler C, Vogt TM, Armstrong GL, Vaughan G, Weltman A, Nainan OV, Dato V, Xia G, Waller K, Amon J, Lee TM, Highbaugh-Battle A, Hembree C, Evenson S, Ruta MA, Williams IT, Fiore AE, Bell BP. An outbreak of hepatitis A associated with green onions. *N Engl J Med* 2005;353:890–897.
12. Donnan EJ, Fielding JE, Gregory JE, Lalor K, Rowe S, Goldsmith P, Antoniou M, Fullerton KE, Knope K, Copland JG, Bowden DS, Tracy SL, Hogg GG, Tan A, Adamopoulos J, Gaston J, Vally H. A multistate outbreak of hepatitis A associated with semidried tomatoes in Australia, 2009. *Clin Infect Dis* 2012;54:775–781.
13. Guillois-Bécel Y, Couturier E, Le Saux JC, Roque-Afonso AM, Le Guyader FS, Le Goas A, Pernès J, Le Behec S, Briand A, Robert C, Dussaix E, Pommepuy M, Vaillant V. An oyster-associated hepatitis A outbreak in France in 2007. *Euro Surveill* 2009;14:pii:19144.

1
2
3 537 14. Tominaga A, Kanda T, Akiike T, Komoda H, Ito K, Abe A, Aruga A, Kaneda S, Saito M, Kiyohara T,
4 538 Wakita T, Ishii K, Yokosuka O, Sugiura N. Hepatitis A outbreak associated with a revolving sushi
5 539 bar in Chiba, Japan: Application of molecular epidemiology. *Hepatol Res* 2012;42:828-834.
6
7
8 540 15. Martin A, Lemon SM. Hepatitis A virus: from discovery to vaccines. *Hepatology* 2006;43:S164–
9 541 172.
10
11 542 16. International Committee on Taxonomy of Viruses. *Virus Taxonomy: 2011 Release (current)*
12 543 [Internet]. [updated 2012; cited 2013 Feb 6]. Available from:
13 544 <http://ictvonline.org/virusTaxonomy.asp?version=2011>.
14
15
16 545 17. Cristina J, Costa-Mattioli M. Genetic variability and molecular evolution of hepatitis A virus. *Virus*
17 546 *Res* 2007;127:151–157.
18
19 547 18. Pérez-Sautu U, Costafreda MI, Caylà J, Tortajada C, Lite J, Bosch A, Pintó RM. Hepatitis a virus
20 548 vaccine escape variants and potential new serotype emergence. *Emerging Infect Dis*
21 549 2011;17:734–737.
22
23
24 550 19. Zhang B, Seitz S, Kusov Y, Zell R, Gauss-Müller V. RNA interaction and cleavage of poly(C)-binding
25 551 protein 2 by hepatitis A virus protease. *Biochem Biophys Res Commun* 2007;364:725–730.
26
27 552 20. Dollenmaier G, Weitz M. Interaction of glyceraldehyde-3-phosphate dehydrogenase with
28 553 secondary and tertiary RNA structural elements of the hepatitis A virus 3' translated and non-
29 554 translated regions. *J Gen Virol* 2003;84:403-414.
30
31
32 555 21. Kaplan G, Totsuka A, Thompson P, Akatsuka T, Moritsugu Y, Feinstone SM. Identification of a
33 556 surface glycoprotein on African green monkey kidney cells as a receptor for hepatitis A virus.
34 557 *EMBO J* 1996;15:4282–4296.
35
36 558 22. Feigelstock D, Thompson P, Mattoo P, Zhang Y, Kaplan GG. The human homolog of HAVcr-1
37 559 codes for a hepatitis A virus cellular receptor. *J Virol*. 1998;72:6621–6628.
38
39
40 560 23. Ichimura T, Bonventre JV, Bailly V, Wei H, Hession CA, Cate RL, Sanicola M. Kidney injury
41 561 molecule-1 (KIM-1), a putative epithelial cell adhesion molecule containing a novel
42 562 immunoglobulin domain, is up-regulated in renal cells after injury. *J Biol Chem* 1998;273:4135–
43 563 4142.
44
45 564 24. Rodriguez-Manzanet R, DeKruyff R, Kuchroo VK, Umetsu DT. The costimulatory role of TIM
46 565 molecules. *Immunol Rev* 2009;229:259–270.
47
48
49 566 25. Sui L, Zhang W, Chen Y, Zheng Y, Wan T, Zhang W, Yang Y, Fang G, Mao J, Cao X. Human
50 567 membrane protein Tim-3 facilitates hepatitis A virus entry into target cells. *Int J Mol Med*
51 568 2006;17:1093–1099.
52
53 569 26. Dotzauer A, Gebhardt U, Bieback K, Göttke U, Kracke A, Mages J, Lemon SM, Vallbracht A.
54 570 Hepatitis A virus-specific immunoglobulin A mediates infection of hepatocytes with hepatitis A
55 571 virus via the asialoglycoprotein receptor. *J Virol* 2000;74:10950–10957.
56
57
58
59
60

27. Tami C, Silberstein E, Manangeeswaran M, Freeman GJ, Umetsu SE, DeKruyff RH, Umetsu DT, Kaplan GG. Immunoglobulin A (IgA) is a natural ligand of hepatitis A virus cellular receptor 1 (HAVCR1), and the association of IgA with HAVCR1 enhances virus-receptor interactions. *J Virol* 2007;81:3437–3446.
28. Dotzauer A, Heitmann A, Laue T, Kraemer L, Schwabe K, Paulmann D, Flehmig B, Vallbracht A. The role of immunoglobulin A in prolonged and relapsing hepatitis A virus infections. *J Gen Virol* 2012;93:754–760.
29. Superti F, Seganti L, Orsi N, Divizia M, Gabrieli R, Panà A. The effect of lipophilic amines on the growth of hepatitis A virus in Frp/3 cells. *Arch Virol* 1987;96:289–296.
30. Superti F, Seganti L, Orsi N, Divizia M, Gabrieli R, Panà A. Effect of cellular function inhibitors on the infection of Frp/3 cells by hepatitis A virus. *Med Microbiol Immunol* 1989;178:29–36.
31. Bishop NE. Examination of potential inhibitors of hepatitis A virus uncoating. *Intervirology* 1998;41:261–271.
32. Bishop NE. Conformational changes in the hepatitis A virus capsid in response to acidic conditions. *J Med Microbiol* 1999;48:443–450.
33. Silberstein E, Xing L, Van de Beek W, Lu J, Cheng H, Kaplan GG. Alteration of hepatitis A virus (HAV) particles by a soluble form of HAV cellular receptor 1 containing the immunoglobulin-and mucin-like regions. *J Virol* 2003;77:8765–8774.
34. Bishop NE, Anderson DA. Uncoating kinetics of hepatitis A virus virions and provirions. *J Virol* 2000;74:3423–3426.
35. Bishop NE. Effect of low pH on the hepatitis A virus maturation cleavage. *Acta Virol* 1999;43:291–296.
36. Bishop NE. Hepatitis A virus replication: an intermediate in the uncoating process. *Intervirology* 2000;43:36–47.
37. Bishop NE, Anderson DA. Early interactions of hepatitis A virus with cultured cells: viral elution and the effect of pH and calcium ions. *Arch Virol* 1997;142:2161–2178.
38. Redondo N, Sanz MA, Steinberger J, Skern T, Kusov Y, Carrasco L. Translation Directed by Hepatitis A Virus IRES in the Absence of Active eIF4F Complex and eIF2. *PLoS ONE* 2012;7:e52065.
39. Borman AM, Kean KM. Intact eukaryotic initiation factor 4G is required for hepatitis A virus internal initiation of translation. *Virology* 1997;237:129–136.
40. Schultheiss T, Kusov YY, Gauss-Müller V. Proteinase 3C of hepatitis A virus (HAV) cleaves the HAV polyprotein P2-P3 at all sites including VP1/2A and 2A/2B. *Virology* 1994;198:275–281.
41. Kusov Y, Gauss-Müller V. Improving proteolytic cleavage at the 3A/3B site of the hepatitis A virus polyprotein impairs processing and particle formation, and the impairment can be complemented in trans by 3AB and 3ABC. *J Virol* 1999;73:9867–9878.

42. De Palma AM, Vliegen I, De Clercq E, Neyts J. Selective inhibitors of picornavirus replication. *Med Res Rev* 2008;28:823–884.
43. Gamarnik AV, Andino R. Switch from translation to RNA replication in a positive-stranded RNA virus. *Genes Dev* 1998;12:2293–2304.
44. Zhang B, Morace G, Gauss-Müller V, Kusov Y. Poly(A) binding protein, C-terminally truncated by the hepatitis A virus proteinase 3C, inhibits viral translation. *Nucleic Acids Res* 2007;35:5975–5984.
45. Schultz DE, Hardin CC, Lemon SM. Specific interaction of glyceraldehyde 3-phosphate dehydrogenase with the 5'-nontranslated RNA of hepatitis A virus. *J Biol Chem* 1996;271:14134–14142.
46. Yi M, Schultz DE, Lemon SM. Functional significance of the interaction of hepatitis A virus RNA with glyceraldehyde 3-phosphate dehydrogenase (GAPDH): opposing effects of GAPDH and polypyrimidine tract binding protein on internal ribosome entry site function. *J Virol* 2000;74:6459–6468.
47. Cordes S, Kusov Y, Heise T, Gauss-Müller V. La autoantigen suppresses IRES-dependent translation of the hepatitis A virus. *Biochem Biophys Res Commun* 2008;368:1014–1019.
48. Teterina NL, Bienz K, Egger D, Gorbalenya AE, Ehrenfeld E. Induction of intracellular membrane rearrangements by HAV proteins 2C and 2BC. *Virology* 1997;237:66–77.
49. Gosert R, Egger D, Bienz K. A cytopathic and a cell culture adapted hepatitis A virus strain differ in cell killing but not in intracellular membrane rearrangements. *Virology* 2000;266:157–169.
50. Yang Y, Liang Y, Qu L, Chen Z, Yi M, Li K, Lemon SM. Disruption of innate immunity due to mitochondrial targeting of a picornaviral protease precursor. *Proc Natl Acad Sci USA* 2007;104:7253–7258.
51. Feng Z, Lemon SM. Hepatitis A virus. In: Ehrenfeld E, Domingo E, Roos RP, editors. *The picornaviruses*. Washington, DC: ASM press; 2010. p383–410.
52. Yang Y, Yi M, Evans DJ, Simmonds P, Lemon SM. Identification of a conserved RNA replication element (cre) within the 3Dpol-coding sequence of hepatoviruses. *J Virol* 2008;82:10118–10128.
53. Ciervo A, Beneduce F, Morace G. Polypeptide 3AB of hepatitis A virus is a transmembrane protein. *Biochem Biophys Res Commun* 1998;249:266–274.
54. Xiang W, Cuconati A, Paul AV, Cao X, Wimmer E. Molecular dissection of the multifunctional poliovirus RNA-binding protein 3AB. *RNA* 1995;1:892–904.
55. Banerjee R, Dasgupta A. Interaction of picornavirus 2C polypeptide with the viral negative-strand RNA. *J Gen Virol* 2001;82:2621–2627.
56. Cohen L, Bénichou D, Martin A. Analysis of deletion mutants indicates that the 2A polypeptide of hepatitis A virus participates in virion morphogenesis. *J Virol* 2002;76:7495–7505.

- 1
2
3 643 57. Martin A, Bénichou D, Chao SF, Cohen LM, Lemon SM. Maturation of the hepatitis A virus capsid
4 644 protein VP1 is not dependent on processing by the 3Cpro proteinase. *J Virol* 1999;73:6220–
5 645 6227.
- 6
7
8 646 58. Rachow A, Gauss-Müller V, Probst C. Homogeneous hepatitis A virus particles. Proteolytic
9 647 release of the assembly signal 2A from procapsids by factor Xa. *J Biol Chem* 2003;278:29744–
10 648 29751.
- 11
12 649 59. Morace G, Kusov Y, Dzagurov G, Beneduce F, Gauss-Muller V. The unique role of domain 2A of
13 650 the hepatitis A virus precursor polypeptide P1-2A in viral morphogenesis. *BMB Rep*
14 651 2008;41:678–683.
- 15
16
17 652 60. Bishop NE, Anderson DA. RNA-dependent cleavage of VP0 capsid protein in provirions of
18 653 hepatitis A virus. *Virology* 1993;197:616–623.
- 19
20 654 61. Tesar M, Jia XY, Summers DF, Ehrenfeld E. Analysis of a potential myristoylation site in hepatitis
21 655 A virus capsid protein VP4. *Virology* 1993;194:616–626.
- 22
23 656 62. Blank CA, Anderson DA, Beard M, Lemon SM. Infection of polarized cultures of human intestinal
24 657 epithelial cells with hepatitis A virus: vectorial release of progeny virions through apical cellular
25 658 membranes. *J Virol* 2000;74:6476–6484.
- 26
27
28 659 63. Snooks MJ, Bhat P, Mackenzie J, Counihan NA, Vaughan N, Anderson DA. Vectorial entry and
29 660 release of hepatitis A virus in polarized human hepatocytes. *J Virol* 2008;82:8733–8742.
- 30
31 661 64. Lemon SM. Type A viral hepatitis. New developments in an old disease. *N Eng J Med*
32 662 1985;313:1059–1067.
- 33
34 663 65. Feinstone SM, Kapikian AZ, Purcell RH. Hepatitis A: detection by immune electron microscopy of
35 664 a viruslike antigen associated with acute illness. *Science* 1973;182:1026–1028.
- 36
37
38 665 66. Mattioli S, Imberti L, Stellini R, Primi D. Mimicry of the immunodominant conformation-
39 666 dependent antigenic site of hepatitis A virus by motifs selected from synthetic peptide libraries.
40 667 *J Virol* 1995;69:5294–5299.
- 41
42 668 67. Sánchez G, Aragonès L, Costafreda MI, Ribes E, Bosch A, Pintó RM. Capsid region involved in
43 669 hepatitis A virus binding to glycoporphin A of the erythrocyte membrane. *J Virol* 2004;78:9807–
44 670 9813.
- 45
46
47 671 68. Costafreda MI, Ribes E, Franch A, Bosch A, Pintó RM. A single mutation in the glycoporphin A
48 672 binding site of hepatitis A virus enhances virus clearance from the blood and results in a lower
49 673 fitness variant. *J Virol* 2012;86:7887–7895.
- 50
51 674 69. Lanford RE, Feng Z, Chavez D, Guerra B, Brasky KM, Zhou Y, Yamane D, Perelson AS, Walker CM,
52 675 Lemon SM. Acute hepatitis A virus infection is associated with a limited type I interferon
53 676 response and persistence of intrahepatic viral RNA. *Proc Natl Acad Sci USA* 2011;108:11223–
54 677 11228.
- 55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

70. Brack K, Berk I, Magulski T, Lederer J, Dotzauer A, Vallbracht A. Hepatitis A virus inhibits cellular antiviral defense mechanisms induced by double-stranded RNA. *J Virol* 2002;76:11920–11930.

71. Fensterl V, Grotheer D, Berk I, Schlemminger S, Vallbracht A, Dotzauer A. Hepatitis A virus suppresses RIG-I-mediated IRF-3 activation to block induction of beta interferon. *J Virol* 2005;79:10968–10977.

72. Paulmann D, Magulski T, Schwarz R, Heitmann L, Flehmig B, Vallbracht A, Dotzauer A. Hepatitis A virus protein 2B suppresses beta interferon (IFN) gene transcription by interfering with IFN regulatory factor 3 activation. *J Gen Virol* 2008;89:1593–1604.

73. Qu L, Feng Z, Yamane D, Liang Y, Lanford RE, Li K, Lemon SM. Disruption of TLR3 signaling due to cleavage of TRIF by the hepatitis A virus protease-polymerase processing intermediate, 3CD. *PLoS Pathog* 2011;7:e1002169.

74. Mukherjee A, Morosky SA, Delorme-Axford E, Dybdahl-Sissoko N, Oberste MS, Wang T, Coyne CB. The coxsackievirus B 3C protease cleaves MAVS and TRIF to attenuate host type I interferon and apoptotic signaling. *PLoS Pathog* 2011;7:e1001311.

75. Lei X, Sun Z, Liu X, Jin Q, He B, Wang J. Cleavage of the adaptor protein TRIF by enterovirus 71 3C inhibits antiviral responses mediated by Toll-like receptor 3. *J Virol* 2011;85:8811–8818.

76. Novoa EM, Ribas de Pouplana L. Speeding with control: codon usage, tRNAs, and ribosomes. *Trends Genet* 2012;28:574-581.

77. Sánchez G, Bosch A, Pintó RM. Genome variability and capsid structural constraints of hepatitis a virus. *J Virol* 2003;77:452–459.

78. Pintó RM, Aragonès L, Costafreda MI, Ribes E, Bosch A. Codon usage and replicative strategies of hepatitis A virus. *Virus Res* 2007;127:158–163.

79. D’Andrea L, Pintó RM, Bosch A, Musto H, Cristina J. A detailed comparative analysis on the overall codon usage patterns in hepatitis A virus. *Virus Res* 2011;157:19–24.

80. Bosch A, Mueller S, Pintó RM. Codon biases and viral fitness. In: Ehrenfeld E, Domingo E, Roos RP, editors. *The picornaviruses*. Washington, DC: ASM press; 2010. p271-283.

81. Marin M. Folding at the rhythm of the rare codon beat. *Biotechnol J* 2008;3:1047–1057.

82. Aragonès L, Guix S, Ribes E, Bosch A, Pintó RM. Fine-tuning translation kinetics selection as the driving force of codon usage bias in the hepatitis A virus capsid. *PLoS Pathog* 2010;6:e1000797.

83. Pintó RM, D’Andrea L, Pérez-Rodríguez FJ, Costafreda MI, Ribes E, Guix S, Bosch A. Hepatitis A virus evolution and the potential emergence of new variants escaping the presently available vaccines. *Future Microbiol* 2012;7:331–346.

84. Brown EA, Zajac AJ, Lemon SM. In vitro characterization of an internal ribosomal entry site (IRES) present within the 5’ nontranslated region of hepatitis A virus RNA: comparison with the IRES of encephalomyocarditis virus. *J Virol* 1994;68:1066–1074.

85. Borman AM, Le Mercier P, Girard M, Kean KM. Comparison of picornaviral IRES-driven internal initiation of translation in cultured cells of different origins. *Nucleic Acids Res* 1997;25:925–932.
86. Manangeeswaran M, Jacques J, Tami C, Konduru K, Amharref N, Perrella O, Casasnovas JM, Umetsu DT, DeKruyff RH, Freeman GJ, Perella A, Kaplan GG. Binding of hepatitis A virus to its cellular receptor 1 inhibits T-regulatory cell functions in humans. *Gastroenterology*. 2012;142:1516–1525.e3.
87. Belkaid Y, Tarbell K. Regulatory T cells in the control of host-microorganism interactions (*). *Annu Rev Immunol* 2009;27:551–589.
88. Jeong S-H, Lee H-S. Hepatitis A: clinical manifestations and management. *Intervirology* 2010;53:15–19.
89. Matricardi PM, Rosmini F, Panetta V, Ferrigno L, Bonini S. Hay fever and asthma in relation to markers of infection in the United States. *J Allergy Clin Immunol* 2002;110:381–387.
90. Linneberg A, Ostergaard C, Tvede M, Andersen LP, Nielsen NH, Madsen F, Frølund L, Dirksen A, Jørgensen T. IgG antibodies against microorganisms and atopic disease in Danish adults: the Copenhagen Allergy Study. *J Allergy Clin Immunol* 2003;111:847–853.
91. Jarvis D, Luczynska C, Chinn S, Burney P. The association of hepatitis A and *Helicobacter pylori* with sensitization to common allergens, asthma and hay fever in a population of young British adults. *Allergy* 2004;59:1063–1067.
92. Gonzalez-Quintela A, Gude F, Boquete O, Aguilera A, Rey J, Meijide LM, Fernandez-Merino MC, Vidal C. Association of hepatitis A virus infection with allergic sensitization in a population with high prevalence of hepatitis A virus exposure. *Allergy* 2005;60:98–103.
93. McIntire JJ, Umetsu SE, Macaubas C, Hoyte EG, Cinnioğlu C, Cavalli-Sforza LL, Barsh GS, Hallmayer JF, Underhill PA, Risch NJ, Freeman GJ, DeKruyff RH, Umetsu DT. Immunology: hepatitis A virus link to atopic disease. *Nature* 2003;425:576.
94. Gao P-S, Mathias RA, Plunkett B, Togias A, Barnes KC, Beaty TH, Huang SK. Genetic variants of the T-cell immunoglobulin mucin 1 but not the T-cell immunoglobulin mucin 3 gene are associated with asthma in an African American population. *J. Allergy Clin Immunol* 2005;115:982–988.
95. Chen J-P, Zhao W-L, He N-H, Gui Q, Xiong J-P, Zhou H-M, Wang Y, Chen S, Zhou P. Association of Hepatitis A exposure and TIM-1 with childhood allergic asthma. *J Asthma* 2012;49:697–702.
96. Kim HY, Eyheramonho MB, Pichavant M, Gonzalez Cambaceres C, Matangkasombut P, Cervio G, Kuperman S, Moreira R, Konduru K, Manangeeswaran M, Freeman GJ, Kaplan GG, DeKruyff RH, Umetsu DT, Rosenzweig SD. A polymorphism in TIM1 is associated with susceptibility to severe hepatitis A virus infection in humans. *J Clin Invest* 2011;121:1111–1118.
97. Binn LN, Lemon SM, Marchwicki RH, Redfield RR, Gates NL, Bancroft WH. Primary isolation and serial passage of hepatitis A virus strains in primate cell cultures. *J Clin Microbiol* 1984;20:28–33.

98. Emerson SU, Huang YK, Purcell RH. 2B and 2C mutations are essential but mutations throughout the genome of HAV contribute to adaptation to cell culture. *Virology* 1993;194:475–480.
99. Yi M, Lemon SM. Replication of subgenomic hepatitis A virus RNAs expressing firefly luciferase is enhanced by mutations associated with adaptation of virus to growth in cultured cells. *J Virol* 2002;76:1171–1180.
100. Day SP, Murphy P, Brown EA, Lemon SM. Mutations within the 5' nontranslated region of hepatitis A virus RNA which enhance replication in BS-C-1 cells. *J Virol* 1992;66:6533–6540.
101. Schultz DE, Honda M, Whetter LE, McKnight KL, Lemon SM. Mutations within the 5' nontranslated RNA of cell culture-adapted hepatitis A virus which enhance cap-independent translation in cultured African green monkey kidney cells. *J Virol* 1996;70:1041–1049.
102. Cohen JI, Rosenblum B, Ticehurst JR, Daemer RJ, Feinstone SM, Purcell RH. Complete nucleotide sequence of an attenuated hepatitis A virus: comparison with wild-type virus. *Proc Natl Acad Sci USA* 1987;84:2497–2501.
103. Taylor KL, Murphy PC, Asher LV, LeDuc JW, Lemon SM. Attenuation phenotype of a cell culture-adapted variant of hepatitis A virus (HM175/p16) in susceptible New World owl monkeys. *J Infect Dis* 1993;168:592–601.
104. Konduru K, Kaplan GG. Stable growth of wild-type hepatitis A virus in cell culture. *J Virol* 2006;80:1352–1360.
105. Lemon SM, Murphy PC, Shields PA, Ping LH, Feinstone SM, Cromeans T, Jansen RW. Antigenic and genetic variation in cytopathic hepatitis A virus variants arising during persistent infection: evidence for genetic recombination. *J Virol* 1991;65:2056–2065.
106. Brack K, Frings W, Dotzauer A, Vallbracht A. A cytopathogenic, apoptosis-inducing variant of hepatitis A virus. *J Virol* 1998;72:3370–3376.
107. Kulka M, Chen A, Ngo D, Bhattacharya SS, Cebula TA, Goswami BB. The cytopathic 18f strain of Hepatitis A virus induces RNA degradation in FrhK4 cells. *Arch Virol* 2003;148:1275–1300.
108. Kulka M, Calvo MS, Ngo DT, Wales SQ, Goswami BB. Activation of the 2-5OAS/RNase L pathway in CVB1 or HAV/18f infected FRhK-4 cells does not require induction of OAS1 or OAS2 expression. *Virology* 2009;388:169–184.
109. Goswami BB, Kulka M, Ngo D, Cebula TA. Apoptosis induced by a cytopathic hepatitis A virus is dependent on caspase activation following ribosomal RNA degradation but occurs in the absence of 2'-5' oligoadenylate synthetase. *Antiviral Res* 2004;63:153–166.
110. Wales SQ, Ngo D, Hida K, Kulka M. Temperature and density dependent induction of a cytopathic effect following infection with non-cytopathic HAV strains. *Virology* 2012;430:30–42.
111. Vallbracht A, Hofmann L, Wurster KG, Flehmig B. Persistent infection of human fibroblasts by hepatitis A virus. *J Gen Virol* 1984;65:609–615.

- 1
2
3 785 112. Crance JM, L  v  que F, Chousterman S, Jouan A, Tr  po C, Deloince R. Antiviral activity of
4 786 recombinant interferon-alpha on hepatitis A virus replication in human liver cells. *Antiviral Res*
5 787 1995;28:69–80.
- 6
7
8 788 113. Yoshiba M, Inoue K, Sekiyama K. Interferon for hepatitis A. *Lancet* 1994;343:288–289.
- 9
10 789 114. Kanda T, Yokosuka O, Imazeki F, Fujiwara K, Nagao K, Saisho H. Amantadine inhibits
11 790 hepatitis A virus internal ribosomal entry site-mediated translation in human hepatoma cells.
12 791 *Biochem Biophys Res Commun* 2005;331:621–629.
- 13
14 792 115. Yang L, Kiyohara T, Kanda T, Imazeki F, Fujiwara K, Gauss-M  ller V, Ishii K, Wakita T,
15 793 Yokosuka O. Inhibitory effects on HAV IRES-mediated translation and replication by a
16 794 combination of amantadine and interferon-alpha. *Viol J* 2010;7:212.
- 17
18
19 795 116. Kanda T, Wu S, Kiyohara T, Nakamoto S, Jiang X, Miyamura T, Imazeki F, Ishii K, Wakita T,
20 796 Yokosuka O. Interleukin-29 suppresses hepatitis A and C viral internal ribosomal entry site-
21 797 mediated translation. *Viral Immunol* 2012;25:379–386.
- 22
23 798 117. Crance JM, Biziagos E, Passagot J, Van Cuyck-Gandr   H, Deloince R. Inhibition of
24 799 hepatitis A virus replication in vitro by antiviral compounds. *J Med Virol* 1990;31:155–160.
- 25
26
27 800 118. PDR Network. Amantadine - Concise Monograph [Internet]. Montvale (NJ): PDR
28 801 Network, LLC; [updated 2013; cited 2013 Feb 6]. Available from:
29 802 <http://www.pdr.net/drugpages/concisemonograph.aspx?concise=411>.
- 30
31 803 119. Siegl G, Eggers HJ. Failure of guanidine and 2-(alpha-hydroxybenzyl)benzimidazole to
32 804 inhibit replication of hepatitis A virus in vitro. *J Gen Virol* 1982;61:111–114.
- 33
34 805 120. Cromeans T, Fields HA, Sobsey MD. Replication kinetics and cytopathic effect of hepatitis
35 806 A virus. *J Gen Virol* 1989;70:2051–2062.
- 36
37
38 807 121. Cho MW, Ehrenfeld E. Rapid completion of the replication cycle of hepatitis A virus
39 808 subsequent to reversal of guanidine inhibition. *Virology* 1991;180:770–780.
- 40
41 809 122. Girond S, Crance JM, Van Cuyck-Gandre H, Renaudet J, Deloince R. Antiviral activity of
42 810 carrageenan on hepatitis A virus replication in cell culture. *Res Virol* 1991;142:261–270.
- 43
44 811 123. Biziagos E, Crance JM, Passagot J, Deloince R. Inhibitory effects of atropine, protamine,
45 812 and their combination on hepatitis A virus replication in PLC/PRF/5 cells. *Antimicrob Agents*
46 813 *Chemother* 1990;34:1112–1117.
- 47
48
49 814 124. Crance JM, L  v  que F, Biziagos E, Van Cuyck-Gandr   H, Jouan A, Deloince R. Studies on
50 815 mechanism of action of glycyrrhizin against hepatitis A virus replication in vitro. *Antiviral Res*
51 816 1994;23:63–76.
- 52
53 817 125. Fiore C, Eisenhut M, Krausse R, Ragazzi E, Pellati D, Armanini D, e.a. Antiviral effects of
54 818 *Glycyrrhiza* species. *Phytother Res* 2008;22:141–148.

- 1
2
3 819 126. Yamamura Y, Kawakami J, Santa T, Kotaki H, Uchino K, Sawada Y, Tanaka N, Iga T.
4 820 Pharmacokinetic profile of glycyrrhizin in healthy volunteers by a new high-performance liquid
5 821 chromatographic method. *J Pharm Sci* 1992;81:1042–1046.
- 7 822 127. Superti F, Seganti L, Orsi N, Divizia M, Gabrieli R, Panà A, Stein ML. Effect of isoflavans
8 823 and isoflavones on the infection of Frp/3 cells by hepatitis A virus. *Antiviral Res* 1989;11:247–
9 824 254.
- 11 825 128. Phillpotts RJ, Wallace J, Tyrrell DA, Freestone DS, Shepherd WM. Failure of oral 4',6-
12 826 dichloroflavan to protect against rhinovirus infection in man. *Arch Virol* 1983;75:115–121.
- 14 827 129. Al-Nakib W, Willman J, Higgins PG, Tyrrell DA, Shepherd WM, Freestone DS. Failure of
15 828 intranasally administered 4', 6-dichloroflavan to protect against rhinovirus infection in man.
16 829 *Arch Virol* 1987;92:255–260.
- 18 830 130. Widell A, Hansson BG, Oberg B, Nordenfelt E. Influence of twenty potentially antiviral
19 831 substances on in vitro multiplication of hepatitis A virus. *Antiviral Res* 1986;6:103–112.
- 21 832 131. Cuyck-Gandré HV, Job A, Burckhart MF, Girond S, Crance JM. Use of digoxigenin-labeled
22 833 RNA probe to test hepatitis A virus antiviral drugs. *Pathol Biol* 1995;43:411–415.
- 24 834 132. Divizia M, Venuti A, Degener AM, Perez-Bercoff R, Panà A. Methisoprinol-effect on the
25 835 replication cycle of human hepatitis A virus. *Microbiologica* 1992;15:323–328.
- 27 836 133. Rashad AA, El-Sabbagh OI, Baraka MM, Ibrahim SM, Pannecouque C, Andrei G, Snoeck R,
28 837 Balzarini J, Mostafa A. Design, synthesis and preliminary antiviral screening of new N-
29 838 phenylpyrazole and dihydroisoxazole derivatives. *Med Chem Res* 2010;19:1025–1035.
- 31 839 134. el-Sabbagh OI, Rady HM. Synthesis of new acridines and hydrazones derived from cyclic
32 840 beta-diketone for cytotoxic and antiviral evaluation. *Eur J Med Chem* 2009;44:3680–3686.
- 34 841 135. Rashad AE, Hegab MI, Abdel-Megeid RE, Micky JA, Abdel-Megeid FME. Synthesis and
35 842 antiviral evaluation of some new pyrazole and fused pyrazolopyrimidine derivatives. *Bioorg Med*
36 843 *Chem* 2008;16:7102–7106.
- 38 844 136. Shamroukh AH, Ali MA. Anti-HAV activity of some newly synthesized triazolo[4,3-
39 845 b]pyridazines. *Arch Pharm (Weinheim)* 2008;341:223–230.
- 41 846 137. Sayed HH, Hashem AI, Yousif NM, El-Sayed WA. Conversion of 3-arylazo-5-phenyl-2(3H)-
42 847 furanones into other heterocycles of anticipated biological activity. *Arch Pharm (Weinheim)*
43 848 2007;340:315–319.
- 45 849 138. Rashad AE, Mohamed MS, Zaki MEA, Fatahala SS. Synthesis and biological evaluation of
46 850 some pyrrolo[2,3-d]pyrimidines. *Arch Pharm (Weinheim)* 2006;339:664–669.
- 48 851 139. Lall MS, Jain RP, Vederas JC. Inhibitors of 3C cysteine proteinases from Picornaviridae.
49 852 *Curr Top Med Chem* 2004;4:1239–1253.
- 51 853 140. Thibaut HJ, De Palma AM, Neyts J. Combating enterovirus replication: state-of-the-art on
52 854 antiviral research. *Biochem Pharmacol* 2012;83:185–192.
- 54
55
56
57
58
59
60

- 1
2
3 855 141. Jewell DA, Swietnicki W, Dunn BM, Malcolm BA. Hepatitis A virus 3C proteinase
4 856 substrate specificity. *Biochemistry* 1992;31:7862–7869.
- 5
6 857 142. Malcolm BA, Lowe C, Shechosky S, McKay RT, Yang CC, Shah VJ, Simon RJ, Vederas JC,
7 858 Santi DV. Peptide aldehyde inhibitors of hepatitis A virus 3C proteinase. *Biochemistry*
8 859 1995;34:8172–8179.
- 9
10
11 860 143. Morris TS, Frormann S, Shechosky S, Lowe C, Lall MS, Gauss-Müller V, Purcell RH,
12 861 Emerson SU, Vederas JC, Malcolm BA. In vitro and ex vivo inhibition of hepatitis A virus 3C
13 862 proteinase by a peptidyl monofluoromethyl ketone. *Bioorg Med Chem* 1997;5:797–807.
- 14
15 863 144. Yin J, Cherney MM, Bergmann EM, Zhang J, Huitema C, Pettersson H, Eltis LD, Vederas
16 864 JC, James MN. An episulfide cation (thiiranium ring) trapped in the active site of HAV 3C
17 865 proteinase inactivated by peptide-based ketone inhibitors. *J Mol Biol* 2006;361:673–686.
- 18
19
20 866 145. Huang Y, Malcolm BA, Vederas JC. Synthesis and testing of azaglutamine derivatives as
21 867 inhibitors of hepatitis A virus (HAV) 3C proteinase. *Bioorg Med Chem* 1999;7:607–619.
- 22
23 868 146. Hill RD, Vederas JC. Azodicarboxamides: A New Class of Cysteine Proteinase Inhibitor for
24 869 Hepatitis A Virus and Human Rhinovirus 3C Enzymes. *J Org Chem* 1999;64:9538–9546.
- 25
26
27 870 147. Lall MS, Karvellas C, Vederas JC. Beta-lactones as a new class of cysteine proteinase
28 871 inhibitors: inhibition of hepatitis A virus 3C proteinase by N-Cbz-serine beta-lactone. *Org Lett*
29 872 1999;1:803–806.
- 30
31 873 148. Lall MS, Ramtohul YK, James MNG, Vederas JC. Serine and threonine beta-lactones: a
32 874 new class of hepatitis A virus 3C cysteine proteinase inhibitors. *J Org Chem* 2002;67:1536–1547.
- 33
34 875 149. Ramtohul YK, Martin NI, Silkin L, James MNG, Vederas JC. Synthesis of pseudoxazolones
35 876 and their inhibition of the 3C cysteine proteinases from hepatitis A virus and human rhinovirus-
36 877 14. *J Chem Soc, Perkin Trans 1* 2002;1351–1359.
- 37
38
39 878 150. Ramtohul YK, James MNG, Vederas JC. Synthesis and evaluation of keto-glutamine
40 879 analogues as inhibitors of hepatitis A virus 3C proteinase. *J Org Chem* 2002;67:3169–3178.
- 41
42 880 151. Huitema C, Zhang J, Yin J, James MNG, Vederas JC, Eltis LD. Heteroaromatic ester
43 881 inhibitors of hepatitis A virus 3C proteinase: Evaluation of mode of action. *Bioorg Med Chem*
44 882 2008;16:5761–5777.
- 45
46
47 883 152. Wang H-M, Liang P-H. Picornaviral 3C protease inhibitors and the dual 3C
48 884 protease/coronaviral 3C-like protease inhibitors. *Expert Opin Ther Pat* 2010;20:59–71.
- 49
50 885 153. Kanda T, Kusov Y, Yokosuka O, Gauss-Müller V. Interference of hepatitis A virus
51 886 replication by small interfering RNAs. *Biochem Biophys Res Commun* 2004;318:341–345.
- 52
53 887 154. Kusov Y, Kanda T, Palmenberg A, Sgro J-Y, Gauss-Müller V. Silencing of hepatitis A virus
54 888 infection by small interfering RNAs. *J Virol* 2006;80:5599–5610.
- 55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

889 155. Kanda T, Zhang B, Kusov Y, Yokosuka O, Gauss-Müller V. Suppression of hepatitis A virus
890 genome translation and replication by siRNAs targeting the internal ribosomal entry site.
891 Biochem Biophys Res Commun 2005;330:1217–1223.

892 156. Debing, Y, Kaplan, GG, Neyts, J, Jochmans, D. Rapid and convenient assays to assess
893 potential inhibitory activity on in vitro hepatitis A replication. Antiviral Res. Forthcoming 2013.

894 157. Van der Schaar HM, Van der Linden L, Lanke KHW, Strating JRPM, Pürstinger G, De Vries
895 E, de Haan CAM, Neyts J, van Kuppeveld FJM. Coxsackievirus mutants that can bypass host
896 factor PI4KIIIβ and the need for high levels of PI4P lipids for replication. Cell Res 2012;22:1576-
897 1592.

898 158. Purcell RH, Emerson SU. Animal models of hepatitis A and E. ILAR J 2001;42:161–177.

899 159. Feigelstock DA, Thompson P, Kaplan GG. Growth of hepatitis A virus in a mouse liver cell
900 line. J Virol 2005;79:2950–2955.

901 160. Hornei B, Kämmerer R, Moubayed P, Frings W, Gauss-Müller V, Dotzauer A.
902 Experimental hepatitis A virus infection in guinea pigs. J Med Virol 2001;64:402–409.

903

10. Figure legends

Figure 1: HAV genome organization and proteolytic processing of the structural proteins. (IRES = internal ribosome entry site, UTR = untranslated region)

Figure 2: Inhibition of IFN β transcription by HAV through proteolytic cleavage of MAVS (by 3ABC) and TRIF (by 3CD) and through direct inhibition of MAVS, IKK ϵ and TBK1 (by 2B). (dsRNA = double-stranded RNA, IKK = inhibitor of NF- κ B kinase, IRF-3 = interferon regulatory factor 3, MAVS = mitochondrial antiviral signaling protein, MDA-5 = melanoma differentiation-associated gene 5, RIG-I = retinoic acid-inducible gene I, TBK1 = TANK-binding kinase 1, TLR3 = toll-like receptor 3, TRIF = Toll/IL-1 receptor domain-containing adaptor inducing IFN β)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

11. Biosketches

Yannick Debing (1988) graduated in 2011 as a pharmacist and master in drug development and is currently a PhD student at the Rega Institute, University of Leuven. His research project addresses the development of novel antiviral strategies against hepatitis A virus and hepatitis E virus.

Hendrik Jan Thibaut (1984) obtained his PhD degree in virology in 2012 at the University of Leuven. He is currently involved in several projects on picornaviruses in the research group of Prof. J. Neyts at the Rega Institute, University of Leuven. His research focuses on the identification and characterization of novel antiviral strategies for the treatment of infections with enteroviruses, with a special interest in protein 2C and the role of the host cell in virus morphogenesis.

Johan Neyts (1966) is full professor of Virology at the University of Leuven, Belgium. His research is focused on the development of novel antiviral strategies against a number of viruses including picornaviruses (such as rhinovirus that cause exacerbations of asthma and COPD), flaviviruses (mainly dengue) and the hepatitis C virus. His work has been published in several book chapters, in more than 280 peer-reviewed papers and in several patents. Three molecules discovered in his lab as HCV inhibitors made it to advanced clinical studies in man for the treatment of chronic HCV infections. He is on the editorial board of a number of journals, member of several national and international scientific committees and advisory boards as well as on the board of the “International Society for Antiviral Research”. He is co-founder and Chief Scientific Officer of Okapi Sciences NV, a company developing antivirals for veterinary use. He teaches medical virology at the school of dentistry and the school of medicine at the University of Leuven. He has been honored with a number of awards including from the International Society for Antiviral Research, the Royal Belgian Academy of Medicine and the Belgian Fund for Scientific Research.




Figure 1: HAV genome organization and proteolytic processing of the structural proteins. (IRES = internal ribosome entry site, UTR = untranslated region)
190x142mm (300 x 300 DPI)




Figure 2: Inhibition of IFNβ transcription by HAV through proteolytic cleavage of MAVS (by 3ABC) and TRIF (by 3CD) and through direct inhibition of MAVS, IKKε and TBK1 (by 2B). (dsRNA = double-stranded RNA, IKK = inhibitor of NF-κB kinase, IRF-3 = interferon regulatory factor 3, MAVS = mitochondrial antiviral signaling protein, MDA-5 = melanoma differentiation-associated gene 5, RIG-I = retinoic acid-inducible gene I, TBK1 = TANK-binding kinase 1, TLR3 = toll-like receptor 3, TRIF = Toll/IL-1 receptor domain-containing adaptor inducing IFNβ)
190x142mm (300 x 300 DPI)

	Antiviral activity	Toxicity	SI	(putative) target	Reference
IFNα-2a	EC ₅₀ : 90 IU/mL	CC ₅₀ : >10000 IU/mL	>100	Induction of antiviral state	112
Amantadine	EC ₅₀ : 58 μ M	CC ₅₀ : 310 μ M	5.3	HAV IRES-mediated translation	117
Guanidine HCl	Variable inhibition between 0.1-3mM	ND	ND	Protein 2C	99, 107, 119-121
Iota-carrageenan	EC ₅₀ : 2.5 μ g/mL	CC ₅₀ : >1000 μ g/mL	>400	Attachment	122
Atropine	84% reduction in viral titer at 1.7mM	CC ₅₀ : >3.5mM	ND	Attachment and/or uncoating	123
Glycyrrhizin	EC ₅₀ : 325 μ M	CC ₅₀ : 5mM	15	Membrane penetration	117
4,6'-dichloroflavan	EC ₅₀ : 6nM	CC ₅₀ : 45 μ M	7500	Entry and/or uncoating	127
4,6'-dichloroisoflavan	EC ₅₀ : 6nM	CC ₅₀ : 90 μ M	15000	Entry and/or uncoating	127
Ribavirin	EC ₅₀ : 94 μ M	CC ₅₀ : 430 μ M	4.6	Inosine 5'- monophosphate dehydrogenase	117
Pyrazofurin	EC ₅₀ : 0.62 μ M	CC ₅₀ : 28 μ M	45	Orotidine 5'-monophosphate decarboxylase	117
Protamine	97% reduction in viral titer at 100 μ g/mL	CC ₅₀ : 200 μ g/mL	ND	RNA polymerase	123
Ac-LAAQ'-fluoromethylketone	96% reduction in viral titer at 5 μ M	ND	ND	Protein 3C ^{pro}	143
Halopyridinyl ester	EC ₅₀ : 53nM	ND	ND	Protein 3C ^{pro}	151

Table 1: Overview of reported inhibitors of HAV with (putative) targets and calculated values for 50% effective concentrations (EC₅₀), 50% inhibitory concentrations (IC₅₀), 50% cytotoxic concentrations (CC₅₀) and selectivity indices (SI) when provided. ND = not determined